

UWThPh-1997-03
 HEPHY-PUB 664/97
 TGU-19
 ITP-SU-97/01
 TU-517
 hep-ph/9701398

QCD corrections to Higgs boson decays into squarks in the Minimal Supersymmetric Standard Model

A. Bartl,¹ H. Eberl,² K. Hidaka,³ T. Kon,⁴
 W. Majerotto² and Y. Yamada⁵

¹*Institut für Theoretische Physik, Universität Wien, A-1090 Vienna, Austria*

²*Institut für Hochenergiephysik der Österreichischen Akademie der Wissenschaften, A-1050 Vienna, Austria*

³*Department of Physics, Tokyo Gakugei University, Koganei, Tokyo 184, Japan*

⁴*Faculty of Engineering, Seikei University, Musashino, Tokyo 180, Japan*

⁵*Department of Physics, Tohoku University, Sendai 980-77, Japan*

Abstract

We calculate the supersymmetric $\mathcal{O}(\alpha_s)$ QCD corrections to the widths of the Higgs boson decays $H^+ \rightarrow t\tilde{b}$ and $H^0, A^0 \rightarrow t\tilde{t}, \tilde{b}\tilde{b}$ in the on-shell scheme within the Minimal Supersymmetric Standard Model. We find that the QCD corrections are significant, but that the squark pair decay modes are still dominant in a wide parameter region.

1 Introduction

It is well known that the Minimal Supersymmetric Standard Model (MSSM) [1] predicts the existence of five physical Higgs bosons h^0, H^0, A^0 , and $H^\pm$ [2, 3]. In order to facilitate experimental Higgs boson searches it is necessary to perform a thorough theoretical study of their decay branching ratios [4]. Their decays into supersymmetric (SUSY) particles can be very important if they are kinematically allowed [5, 6, 7, 8]. The decays into the 3rd-generation squarks $\tilde{t}$ and $\tilde{b}$ can play a special rôle because they can be much lighter than the other squarks and the decays can be strongly enhanced due to their large Yukawa couplings and their large $\tilde{q}_L - \tilde{q}_R$ mixings [6, 7, 8]. The tree-level results of Refs. [6, 8] show that the decay modes $H^+ \rightarrow \tilde{t}\tilde{b}$ and $H^0, A^0 \rightarrow \tilde{t}\tilde{t}, \tilde{b}\tilde{b}$ can be dominant in a large region of the parameter space of the MSSM, and that this could have an important impact on searches for H^+, H^0 , and A^0 at future colliders.

The SUSY QCD corrections to the decays $H^+ \rightarrow \tilde{t}\tilde{b}$ and $H^0, A^0 \rightarrow \tilde{t}\tilde{t}, \tilde{b}\tilde{b}$ can be large [9]. This suggests that the QCD corrections to $H^+ \rightarrow \tilde{t}\tilde{b}$ and $H^0, A^0 \rightarrow \tilde{t}\tilde{t}, \tilde{b}\tilde{b}$ could also be large. Therefore it is very important to examine whether the results of Refs. [6, 8] remain valid after including the QCD corrections. In Ref. [10] it was shown that the QCD corrections to $H^+ \rightarrow \tilde{t}\tilde{b}$ can be significant in the $\overline{\text{DR}}$ renormalization scheme, but that they do not invalidate the result in Ref. [6] on the dominance of the $H^+ \rightarrow \tilde{t}\tilde{b}$ mode in a large parameter region.

In the present paper we extend our study to the decays of the charged and neutral Higgs bosons. We use the on-shell scheme which is more appropriate for the discussion of physical observables. We calculate the complete $\mathcal{O}(\alpha_s)$ QCD corrections to the widths of the decays $H^+ \rightarrow \tilde{t}\tilde{b}$ and $H^0, A^0 \rightarrow \tilde{t}\tilde{t}, \tilde{b}\tilde{b}$ within the MSSM including all quark mass terms and $\tilde{q}_L - \tilde{q}_R$ mixings. The main complication here is that the $\tilde{q}_L - \tilde{q}_R$ mixing angles are renormalized by the SUSY-QCD corrections. This problem was first solved in Ref. [11] in the treatment of $e^+e^- \rightarrow \tilde{q}\tilde{q}$, where a suitable renormalization condition for the squark-mixing angle was found. The method was also applied in [12, 13, 14, 15] to $\tilde{q}_i \rightarrow q\tilde{\chi}_j^{0,\pm}$ and $t \rightarrow \tilde{t}_1\tilde{\chi}_1^0$. In the present study we use the renormalization prescription as given in [11]. Furthermore, we point out that special attention must be paid to the soft SUSY-breaking parameter $M_{\tilde{Q}}$, which enters the stop and sbottom mass matrices: in the on-shell scheme, the renormalized $M_{\tilde{Q}}$ in the stop sector is different from that in the sbottom sector.

We find that the QCD corrections to the squark pair widths are significant, but that the squark pair modes ($H^+ \rightarrow \tilde{t}\tilde{b}$ and $H^0, A^0 \rightarrow \tilde{t}\tilde{t}, \tilde{b}\tilde{b}$) are still dominant in a wide parameter range.

2 Tree level result

We first review the tree level results [6, 8]. The squark mass matrix in the basis $(\tilde{q}_L, \tilde{q}_R)$, with $\tilde{q} = \tilde{t}$ or $\tilde{b}$, is given by [2, 3]

$$\begin{pmatrix} m_{LL}^2 & m_{LR}^2 \\ m_{RL}^2 & m_{RR}^2 \end{pmatrix} = (R^{\tilde{q}})^\dagger \begin{pmatrix} m_{\tilde{q}_1}^2 & 0 \\ 0 & m_{\tilde{q}_2}^2 \end{pmatrix} R^{\tilde{q}}, \quad (1)$$

where

$$m_{LL}^2 = M_Q^2 + m_q^2 + m_Z^2 \cos 2\beta (I_q^{3L} - e_q \sin^2 \theta_W), \quad (2)$$

$$m_{RR}^2 = M_{\{\tilde{U}, \tilde{D}\}}^2 + m_q^2 + m_Z^2 \cos 2\beta e_q \sin^2 \theta_W, \quad (3)$$

$$m_{LR}^2 = m_{RL}^2 = \begin{cases} m_t(A_t - \mu \cot \beta) & (\tilde{q} = \tilde{t}) \\ m_b(A_b - \mu \tan \beta) & (\tilde{q} = \tilde{b}) \end{cases}, \quad (4)$$

and

$$R_{i\alpha}^{\tilde{q}} = \begin{pmatrix} \cos \theta_{\tilde{q}} & \sin \theta_{\tilde{q}} \\ -\sin \theta_{\tilde{q}} & \cos \theta_{\tilde{q}} \end{pmatrix}. \quad (5)$$

The mass eigenstates $\tilde{q}_i (i = 1, 2)$ (with $m_{\tilde{q}_1} < m_{\tilde{q}_2}$) are related to the $SU(2)_L$ eigenstates $\tilde{q}_\alpha (\alpha = L, R)$ by $\tilde{q}_i = R_{i\alpha}^{\tilde{q}} \tilde{q}_\alpha$.

The tree-level decay width of $H^k \rightarrow \tilde{q}_i \tilde{q}_j^*$ is then given by (see Fig. 1a)

$$\Gamma^{tree}(H^k \rightarrow \tilde{q}_i \tilde{q}_j^*) = \frac{N_C \kappa}{16\pi m_{H^k}^3} |G_{ijk}^{\tilde{q}}|^2. \quad (6)$$

For $k = 1, 2, 3$ H^k denotes the neutral Higgs bosons (i. e. $H^1 \equiv h^0$, $H^2 \equiv H^0$, $H^3 \equiv A^0$) and $\tilde{q} = \tilde{t}, \tilde{b}$. For $k = 4$ one has $H^4 \equiv H^\pm$ and $\tilde{q}_i \equiv \tilde{t}_i, \tilde{q}_j \equiv \tilde{b}_j$, and the upper index $\tilde{q}$ in $G_{ijk}^{\tilde{q}}$ is omitted. $\kappa = \kappa(m_{H^k}^2, m_{\tilde{q}_i}^2, m_{\tilde{q}_j}^2)$, $\kappa(x, y, z) \equiv ((x - y - z)^2 - 4yz)^{1/2}$, $N_C = 3$, and the $H^k \tilde{q}_i^* \tilde{q}_j$ couplings [2, 3] are given by

$$G_{ij2}^{\tilde{t}} \equiv G(H^0 \tilde{t}_i^* \tilde{t}_j) = \left(R^{\tilde{t}} G_{LR}^{\tilde{t}} (R^{\tilde{t}})^T \right)_{ij} = -g R^{\tilde{t}} \begin{pmatrix} \frac{m_Z}{c_W} (I_t^{3L} - e_t s_W^2) c_{\alpha+\beta} + \frac{m_t^2}{m_W s_\beta} s_\alpha & \frac{m_t}{2m_W s_\beta} (A_t s_\alpha - \mu c_\alpha) \\ \frac{m_t}{2m_W s_\beta} (A_t s_\alpha - \mu c_\alpha) & \frac{m_Z}{c_W} e_t s_W^2 c_{\alpha+\beta} + \frac{m_t^2}{m_W s_\beta} s_\alpha \end{pmatrix} (R^{\tilde{t}})^T, \quad (7)$$

$$G_{ij2}^{\tilde{b}} \equiv G(H^0 \tilde{b}_i^* \tilde{b}_j) = \left(R^{\tilde{b}} G_{LR}^{\tilde{b}} (R^{\tilde{b}})^T \right)_{ij} = -g R^{\tilde{b}} \begin{pmatrix} \frac{m_Z}{c_W} (I_b^{3L} - e_b s_W^2) c_{\alpha+\beta} + \frac{m_b^2}{m_W c_\beta} c_\alpha & \frac{m_b}{2m_W c_\beta} (A_b c_\alpha - \mu s_\alpha) \\ \frac{m_b}{2m_W c_\beta} (A_b c_\alpha - \mu s_\alpha) & \frac{m_Z}{c_W} e_b s_W^2 c_{\alpha+\beta} + \frac{m_b^2}{m_W c_\beta} c_\alpha \end{pmatrix} (R^{\tilde{b}})^T, \quad (8)$$

$$G_{ij1}^{\tilde{q}} \equiv G(h^0 \tilde{q}_i^* \tilde{q}_j) = \{G(H^0 \tilde{q}_i^* \tilde{q}_j) \text{ with } \alpha \rightarrow \alpha + \frac{1}{2}\pi, \text{ (i. e. } \sin \alpha \equiv s_\alpha \rightarrow c_\alpha, \cos \alpha \equiv c_\alpha \rightarrow -s_\alpha, \text{ and } \cos(\alpha + \beta) \equiv c_{\alpha+\beta} \rightarrow -s_{\alpha+\beta})\}, \quad (9)$$

$$G_{ij3}^{\tilde{t}} = G(A^0 \tilde{t}_i^* \tilde{t}_j) = \frac{ig}{2m_W} \begin{pmatrix} 0 & m_t(A_t \cot \beta + \mu) \\ -m_t(A_t \cot \beta + \mu) & 0 \end{pmatrix}, \quad (10)$$

$$G_{ij3}^{\tilde{b}} = G(A^0 \tilde{b}_i^* \tilde{b}_j) = \frac{ig}{2m_W} \begin{pmatrix} 0 & m_b(A_b \tan \beta + \mu) \\ -m_b(A_b \tan \beta + \mu) & 0 \end{pmatrix}, \quad (11)$$

$$G_{ij4} \equiv G(H^+ \tilde{t}_i^* \tilde{b}_j) = \frac{g}{\sqrt{2}m_W} R^{\tilde{t}} \begin{pmatrix} m_b^2 \tan \beta + m_t^2 \cot \beta - m_W^2 \sin 2\beta & m_b(A_b \tan \beta + \mu) \\ m_t(A_t \cot \beta + \mu) & 2m_t m_b / \sin 2\beta \end{pmatrix} (R^{\tilde{b}})^T. \quad (12)$$

Here g is the SU(2) coupling, α is the mixing angle in the CP even neutral Higgs boson sector, $c_W \equiv \cos \theta_W$ and $s_W \equiv \sin \theta_W$.

3 QCD corrections

The $\mathcal{O}(\alpha_s)$ QCD virtual corrections to $H^k \rightarrow \tilde{q}_i \tilde{\bar{q}}_j$ stem from the diagrams of Fig. 1b (vertex corrections) and Fig. 1c (wave-function corrections). All parameters of the QCD interacting particles, appearing in the tree-level mass matrix of eqs. (1) – (5) and the tree-level couplings of eqs. (7) – (12), have to be renormalized. These are the soft-SUSY-breaking squark masses $M_{\tilde{Q}, \tilde{U}, \tilde{D}}$, the quark masses $m_{t,b}$, the trilinear couplings $A_{t,b}$, the squark masses $m_{\tilde{q}_{1,2}}$, and the mixing angles $\theta_{\tilde{t}, \tilde{b}}$ (α , β , and μ are of course not renormalized by QCD). In this paper we use the on-shell renormalization scheme.

The one-loop corrected decay amplitudes $G_{ijk}^{\tilde{q} \text{corr}}$ can be expressed as

$$G_{ijk}^{\tilde{q} \text{corr}} = G_{ijk}^{\tilde{q}} + \delta G_{ijk}^{\tilde{q}(v)} + \delta G_{ijk}^{\tilde{q}(w)} + \delta G_{ijk}^{\tilde{q}(0)}, \quad (13)$$

where $G_{ijk}^{\tilde{q}}$ are defined by eqs. (7) – (12) in terms of the on-shell parameters, and $\delta G_{ijk}^{\tilde{q}(v)}$ and $\delta G_{ijk}^{\tilde{q}(w)}$ are the vertex and squark wave-function corrections, respectively. $\delta G_{ijk}^{\tilde{q}(0)}$ denotes the counterterm caused by the on-shell renormalization. They get contributions from the gluon, gluino and squark exchange. (Again the upper index $\tilde{q}$ has to be omitted for $k = 4$.)

The gluon exchange contributions to the vertex corrections are (see Fig. 1b)

$$\begin{aligned} \delta G_{ijk}^{\tilde{q}(v,g)} &= \frac{\alpha_s C_F}{4\pi} G_{ijk}^{\tilde{q}} \left[B_0(m_{\tilde{q}_i}^2, 0, m_{\tilde{q}_i}^2) + B_0(m_{\tilde{q}_j}^2, 0, m_{\tilde{q}_j}^2) - B_0(m_{H^k}^2, m_{\tilde{q}_i}^2, m_{\tilde{q}_j}^2) \right. \\ &\quad \left. - 2(m_{H^k}^2 - m_{\tilde{q}_i}^2 - m_{\tilde{q}_j}^2) C_0(\lambda^2, m_{\tilde{q}_i}^2, m_{\tilde{q}_j}^2) \right]. \end{aligned} \quad (14)$$

The gluino exchange contributions to them are (see Fig. 1b)

$$\begin{aligned}\delta G_{ijk}^{\tilde{q}(v,\tilde{g})} &= \frac{\alpha_s C_F}{2\pi} s_k^q [2\delta_{ij} m_q B_0(m_{H^k}^2, m_q^2, m_q^2) \\ &\quad + \{\delta_{ij} m_q + S_{ij}^{\tilde{q}} m_{\tilde{g}}\} (B_0(m_{\tilde{q}_i}^2, m_{\tilde{g}}^2, m_q^2) + B_0(m_{\tilde{q}_j}^2, m_{\tilde{g}}^2, m_q^2)) + \{ (4m_q^2 - \\ &\quad m_{H^k}^2) S_{ij}^{\tilde{q}} m_{\tilde{g}} + (2m_q^2 + 2m_{\tilde{g}}^2 - m_{\tilde{q}_i}^2 - m_{\tilde{q}_j}^2) \delta_{ij} m_q \} C_0(m_{\tilde{g}}^2, m_q^2, m_q^2)] , \quad (15)\end{aligned}$$

for $k = 1, 2$,

$$\begin{aligned}\delta G_{ij3}^{\tilde{q}(v,\tilde{g})} &= -\frac{\alpha_s C_F}{2\pi} a^q [A_{ij}^{\tilde{q}} m_q (B_0(m_{\tilde{q}_i}^2, m_{\tilde{g}}^2, m_q^2) - B_0(m_{\tilde{q}_j}^2, m_{\tilde{g}}^2, m_q^2)) \\ &\quad + \epsilon_{ij} m_{\tilde{g}} (B_0(m_{\tilde{q}_i}^2, m_{\tilde{g}}^2, m_q^2) + B_0(m_{\tilde{q}_j}^2, m_{\tilde{g}}^2, m_q^2)) \\ &\quad + \{ (m_{\tilde{q}_i}^2 - m_{\tilde{q}_j}^2) A_{ij}^{\tilde{q}} m_q - m_{H^k}^2 \epsilon_{ij} m_{\tilde{g}} \} C_0(m_{\tilde{g}}^2, m_q^2, m_q^2)] , \quad (16)\end{aligned}$$

and

$$\begin{aligned}\delta G_{ij4}^{(v,\tilde{g})} &= \frac{\alpha_s C_F}{2\pi} [\{ (\alpha_{LL})_{ij} (m_t y_2 + m_b y_1) + (\alpha_{RR})_{ij} (m_t y_1 + m_b y_2) \} B_0(m_H^2, m_b^2, m_t^2) \\ &\quad + \{ (\alpha_{LL})_{ij} m_t y_2 + (\alpha_{LR})_{ij} m_{\tilde{g}} y_1 + (\alpha_{RL})_{ij} m_{\tilde{g}} y_2 + (\alpha_{RR})_{ij} m_t y_1 \} B_0(m_{\tilde{t}_i}^2, m_{\tilde{g}}^2, m_t^2) \\ &\quad + \{ (\alpha_{LL})_{ij} m_b y_1 + (\alpha_{LR})_{ij} m_{\tilde{g}} y_1 + (\alpha_{RL})_{ij} m_{\tilde{g}} y_2 + (\alpha_{RR})_{ij} m_b y_2 \} B_0(m_{\tilde{b}_j}^2, m_{\tilde{g}}^2, m_b^2) \\ &\quad + \{ (m_t^2 + m_b^2 - m_H^2) m_{\tilde{g}} ((\alpha_{LR})_{ij} y_1 + (\alpha_{RL})_{ij} y_2) \\ &\quad + (m_b^2 + m_{\tilde{g}}^2 - m_{\tilde{b}_j}^2) m_t ((\alpha_{LL})_{ij} y_2 + (\alpha_{RR})_{ij} y_1) \\ &\quad + (m_t^2 + m_{\tilde{g}}^2 - m_{\tilde{t}_i}^2) m_b ((\alpha_{LL})_{ij} y_1 + (\alpha_{RR})_{ij} y_2) \\ &\quad + 2m_{\tilde{g}} m_t m_b ((\alpha_{LR})_{ij} y_2 + (\alpha_{RL})_{ij} y_1) \} C_0(m_{\tilde{g}}^2, m_t^2, m_b^2)] . \quad (17)\end{aligned}$$

The vertex corrections due to the four-squark interaction are (see Fig. 1b)

$$\delta G_{ijk}^{\tilde{q}(v,\tilde{q})} = -\frac{\alpha_s C_F}{4\pi} \sum_{n,l} B_0(m_{H^k}^2, m_{\tilde{q}_n}^2, m_{\tilde{q}_l}^2) G_{nlk}^{\tilde{q}} A_{in}^{\tilde{q}} A_{lj}^{\tilde{q}} , \quad (k = 1, 2, 3), \quad (18)$$

$$\delta G_{ij4}^{(v,\tilde{q})} = -\frac{\alpha_s C_F}{4\pi} \sum_{n,l} B_0(m_{H^+}^2, m_{\tilde{t}_n}^2, m_{\tilde{b}_l}^2) G_{nl4} A_{in}^{\tilde{t}} A_{lj}^{\tilde{b}} . \quad (19)$$

s_k^q ($k = 1, 2$), a^q , y_1 , and y_2 in eqs. (15), (16), and (17) are the Yukawa couplings [2]:

$$\mathcal{L} = s_1^q h^0 \bar{q} q + s_2^q H^0 \bar{q} q + a_q A^0 \bar{q} \gamma^5 q + H^+ \bar{t} (y_1 P_R + y_2 P_L) b , \quad (20)$$

with

$$\begin{aligned}s_1^t &= -g \frac{m_t \cos \alpha}{2m_W \sin \beta} , & s_1^b &= g \frac{m_b \sin \alpha}{2m_W \cos \beta} , \\ s_2^t &= -g \frac{m_t \sin \alpha}{2m_W \sin \beta} , & s_2^b &= -g \frac{m_b \cos \alpha}{2m_W \cos \beta} , \\ a^t &= ig \frac{m_t \cot \beta}{2m_W} , & a^b &= ig \frac{m_b \tan \beta}{2m_W} , \\ y_1 &= \frac{g}{\sqrt{2}m_W} m_b \tan \beta = h_b \sin \beta \quad \text{and} \quad y_2 = \frac{g}{\sqrt{2}m_W} m_t \cot \beta = h_t \cos \beta .\end{aligned} \quad (21)$$

$C_F = 4/3$, δ_{ij} is the unit matrix, ϵ_{ij} is totally antisymmetric with $\epsilon_{12} = 1$,

$$A^{\tilde{q}} = \begin{pmatrix} \cos 2\theta_{\tilde{q}} & -\sin 2\theta_{\tilde{q}} \\ -\sin 2\theta_{\tilde{q}} & -\cos 2\theta_{\tilde{q}} \end{pmatrix}, \quad S^{\tilde{q}} = \begin{pmatrix} -\sin 2\theta_{\tilde{q}} & -\cos 2\theta_{\tilde{q}} \\ -\cos 2\theta_{\tilde{q}} & \sin 2\theta_{\tilde{q}} \end{pmatrix} = \epsilon_{ik} A_{kj}^{\tilde{q}}, \quad (22)$$

$$(\alpha_{\sigma\rho})_{ij} = (2\delta_{\sigma\rho} - 1) R_{i\sigma}^{\tilde{t}} R_{j\rho}^{\tilde{b}}, \quad (23)$$

and $m_{\tilde{g}}$ is the gluino mass. A gluon mass λ is introduced to regularize the infrared divergences. The UV divergences are regularized by dimensional reduction (DR) [16, 17] which preserves supersymmetry at least at one-loop order. We use the usual one-, two-, and three-point functions A_0 , B_0 , B_1 , and C_0 [18]:

$$\begin{aligned} A_0(m^2) &= \int \frac{d^D q}{i\pi^2} \frac{1}{q^2 - m^2}, \\ [B_0, k^\mu B_1](k^2, m_1^2, m_2^2) &= \int \frac{d^D q}{i\pi^2} \frac{[1, q^\mu]}{(q^2 - m_1^2)((q+k)^2 - m_2^2)}, \\ C_0(m_0^2, m_1^2, m_2^2) &= \int \frac{d^D q}{i\pi^2} \frac{1}{(q^2 - m_0^2)((q+k_{\tilde{q}_i})^2 - m_1^2)((q+k_{\tilde{q}_j})^2 - m_2^2)}. \end{aligned}$$

In these integrals $k_{\tilde{q}_i}$ and $-k_{\tilde{q}_j}$ are the external momenta of $\tilde{q}_i$ and $\tilde{q}_j$, respectively.

The squark wave-function corrections $\delta G_{ijk}^{\tilde{q}(w)}$ can be expressed as

$$\delta G_{ijk}^{\tilde{q}(w)} = -\frac{1}{2} [\dot{\Pi}_{ii}^{\tilde{q}}(m_{\tilde{q}_i}^2) + \dot{\Pi}_{jj}^{\tilde{q}}(m_{\tilde{q}_j}^2)] G_{ijk}^{\tilde{q}} - \frac{\Pi_{ii'}^{\tilde{q}}(m_{\tilde{q}_i}^2)}{m_{\tilde{q}_i}^2 - m_{\tilde{q}_{i'}}^2} G_{i'jk}^{\tilde{q}} - \frac{\Pi_{jj'}^{\tilde{q}}(m_{\tilde{q}_j}^2)}{m_{\tilde{q}_j}^2 - m_{\tilde{q}_{j'}}^2} G_{ij'k}^{\tilde{q}}. \quad (24)$$

Here and in the following $i \neq i'$ and $j \neq j'$. $\Pi_{ij}^{\tilde{q}}(k^2)$ are the one-loop corrections to the two-point functions of $\tilde{q}_i \tilde{q}_j$, which are obtained from the graphs of Fig. 1c. $\dot{\Pi}(k^2)$ denotes the derivative with respect to k^2 . The last two terms in (24) represent the corrections due to squark mixing. Note that for H^+ decay ($k = 4$) the subscripts i and i' are attached to $\tilde{t}$ and j and j' to $\tilde{b}$. The explicit forms of the self-energies and their derivatives of the diagonal parts are

$$\Pi_{ij}^{\tilde{q}(g)}(k^2) = -\frac{\alpha_s C_F}{4\pi} \delta_{ij} [(3k^2 + m_{\tilde{q}_i}^2) B_0(k^2, \lambda^2, m_{\tilde{q}_i}^2) + 2k^2 B_1(k^2, \lambda^2, m_{\tilde{q}_i}^2)], \quad (25)$$

$$\begin{aligned} \Pi_{ij}^{\tilde{q}(\tilde{g})}(k^2) &= -\frac{\alpha_s C_F}{\pi} [(A_0(m_q^2) + k^2 B_1(k^2, m_{\tilde{g}}^2, m_q^2)) \delta_{ij} \\ &\quad + (m_{\tilde{g}}^2 \delta_{ij} + (S^{\tilde{q}})_{ij} m_q m_{\tilde{g}}) B_0(k^2, m_{\tilde{g}}^2, m_q^2)], \end{aligned} \quad (26)$$

$$\Pi_{ij}^{\tilde{q}(\tilde{q})}(k^2) = \frac{\alpha_s C_F}{4\pi} \begin{pmatrix} \cos^2 2\theta_{\tilde{q}} A_0(m_{\tilde{q}_1}^2) + \sin^2 2\theta_{\tilde{q}} A_0(m_{\tilde{q}_2}^2) & \frac{1}{2} \sin 4\theta_{\tilde{q}} (A_0(m_{\tilde{q}_2}^2) - A_0(m_{\tilde{q}_1}^2)) \\ \frac{1}{2} \sin 4\theta_{\tilde{q}} (A_0(m_{\tilde{q}_2}^2) - A_0(m_{\tilde{q}_1}^2)) & \sin^2 2\theta_{\tilde{q}} A_0(m_{\tilde{q}_1}^2) + \cos^2 2\theta_{\tilde{q}} A_0(m_{\tilde{q}_2}^2) \end{pmatrix}, \quad (27)$$

$$\begin{aligned}\dot{\Pi}_{ii}^{\tilde{q}(g)}(m_{\tilde{q}_i}^2) &= -\frac{\alpha_s C_F}{4\pi} \left[3B_0(m_{\tilde{q}_i}^2, 0, m_{\tilde{q}_i}^2) + 2B_1(m_{\tilde{q}_i}^2, 0, m_{\tilde{q}_i}^2) \right. \\ &\quad \left. + 4m_{\tilde{q}_i}^2 \dot{B}_0(m_{\tilde{q}_i}^2, \lambda^2, m_{\tilde{q}_i}^2) + 2m_{\tilde{q}_i}^2 \dot{B}_1(m_{\tilde{q}_i}^2, 0, m_{\tilde{q}_i}^2) \right],\end{aligned}\quad (28)$$

$$\begin{aligned}\dot{\Pi}_{ii}^{\tilde{q}(\tilde{g})}(m_{\tilde{q}_i}^2) &= -\frac{\alpha_s C_F}{\pi} \left[m_{\tilde{g}}^2 \dot{B}_0(m_{\tilde{q}_i}^2, m_{\tilde{g}}^2, m_q^2) + B_1(m_{\tilde{q}_i}^2, m_{\tilde{g}}^2, m_q^2) \right. \\ &\quad \left. + m_{\tilde{q}_i}^2 \dot{B}_1(m_{\tilde{q}_i}^2, m_{\tilde{g}}^2, m_q^2) + (-)^i \sin 2\theta_{\tilde{q}} m_q m_{\tilde{g}} \dot{B}_0(m_{\tilde{q}_i}^2, m_{\tilde{g}}^2, m_q^2) \right],\end{aligned}\quad (29)$$

$$\dot{\Pi}_{ii}^{\tilde{q}(\tilde{q})}(m_{\tilde{q}_i}^2) = 0. \quad (30)$$

Now we discuss the shifts $\delta G_{ij2}^{\tilde{q}(0)}$ in eq. (13). From eqs. (7) and (8) it follows:

$$\delta G_{ij2}^{\tilde{q}(0)} = \left(R^{\tilde{q}} \delta G_{LR}^{\tilde{q}} (R^{\tilde{q}})^T + \delta R^{\tilde{q}} G_{LR}^{\tilde{q}} (R^{\tilde{q}})^T + R^{\tilde{q}} G_{LR}^{\tilde{q}} \delta (R^{\tilde{q}})^T \right)_{ij}. \quad (31)$$

Using eq. (5) one further gets:

$$\delta G_{ij2}^{\tilde{q}(0)} = (R^{\tilde{q}})_{ik} (\delta G_{LR}^{\tilde{q}})_{kl} (R^{\tilde{q}})_{jl} - \left((-1)^i G_{i'j2}^{\tilde{q}} + (-1)^j G_{ij'2}^{\tilde{q}} \right) \delta \theta_{\tilde{q}}. \quad (32)$$

From eqs. (7) and (8) it directly follows:

$$\delta G_{LR}^{\tilde{t}} = -\frac{g}{2m_W s_\beta} \begin{pmatrix} 4m_t s_\alpha \delta m_t & \delta(m_t A_t) s_\alpha - \mu c_\alpha \delta m_t \\ \delta(m_t A_t) s_\alpha - \mu c_\alpha \delta m_t & 4m_t s_\alpha \delta m_t \end{pmatrix}, \quad (33)$$

$$\delta G_{LR}^{\tilde{b}} = -\frac{g}{2m_W c_\beta} \begin{pmatrix} 4m_b c_\alpha \delta m_b & \delta(m_b A_b) c_\alpha - \mu s_\alpha \delta m_b \\ \delta(m_b A_b) c_\alpha - \mu s_\alpha \delta m_b & 4m_b c_\alpha \delta m_b \end{pmatrix}. \quad (34)$$

To get the correction terms $\delta G_{ij1}^{\tilde{q}(0)}$ one makes the same replacements as in eq. (9):

$$\delta G_{ij1}^{\tilde{q}(0)} = \left(\delta G_{ij2}^{\tilde{q}(0)} \text{ with } \alpha \rightarrow \alpha + \frac{1}{2}\pi \right). \quad (35)$$

For the couplings to the A^0 boson, eqs. (10) and (11) one gets the correction terms

$$\delta G_{ij3}^{\tilde{q}(0)} = \frac{ig}{2m_W} \begin{pmatrix} 0 & \delta(m_q A_q) \{ \cot \beta, \tan \beta \} + \mu \delta m_q \\ -(\delta(m_q A_q) \{ \cot \beta, \tan \beta \} + \mu \delta m_q) & 0 \end{pmatrix}, \quad (36)$$

where $\cot \beta$ ($\tan \beta$) has to be taken for $\tilde{q} = \tilde{t}$ ($\tilde{b}$).

For the $H^+ \rightarrow \tilde{t} \tilde{b}$ ($k = 4$) we have

$$\delta G_{ij4}^{(0)} = \left(R^{\tilde{t}} \delta G_{LR}^{\tilde{b}} (R^{\tilde{b}})^T \right)_{ij} - (-1)^i G_{i'j4} \delta \theta_{\tilde{t}} - (-1)^j G_{ij'4} \delta \theta_{\tilde{b}}, \quad (37)$$

and

$$\begin{aligned}\delta G_{LR} &= \\ \frac{g}{\sqrt{2}m_W} &\begin{pmatrix} 2m_b \delta m_b \tan \beta + 2m_t \delta m_t \cot \beta & \delta(m_b A_b) \tan \beta + \delta m_b \mu \\ \delta(m_t A_t) \cot \beta + \delta m_t \mu & 2(\delta m_t m_b + m_t \delta m_b) / \sin 2\beta \end{pmatrix}. \end{aligned}\quad (38)$$

We now give the formulae for δm_q , $\delta(m_q A_q)$, and $\delta\theta_{\tilde{q}}$ in the on-shell scheme. These terms consist of three parts, denoted by the superscripts g , $\tilde{g}$, and $\bar{q}$. Therefore we can write $\delta G_{ijk}^{\tilde{q}(0)} \equiv \delta G_{ijk}^{\tilde{q}(0,g)} + \delta G_{ijk}^{\tilde{q}(0,\tilde{g})} + \delta G_{ijk}^{\tilde{q}(0,\bar{q})}$. Since the renormalized m_q is taken to be the pole mass, one gets

$$\begin{aligned}\delta m_q &= \delta m_q^{(g)} + \delta m_q^{(\tilde{g})}, \\ \delta m_q^{(g)} &= -\frac{\alpha_s C_F}{2\pi} m_q (B_0(m_q^2, 0, m_q^2) - B_1(m_q^2, 0, m_q^2)), \text{ and} \\ \delta m_q^{(\tilde{g})} &= -\frac{\alpha_s C_F}{4\pi} [\sin 2\theta_{\tilde{q}} m_{\tilde{g}} (B_0(m_q^2, m_{\tilde{g}}^2, m_{\tilde{q}_1}^2) - B_0(m_q^2, m_{\tilde{g}}^2, m_{\tilde{q}_2}^2)) \\ &\quad + m_q (B_1(m_q^2, m_{\tilde{g}}^2, m_{\tilde{q}_1}^2) + B_1(m_q^2, m_{\tilde{g}}^2, m_{\tilde{q}_2}^2))] ,\end{aligned}\quad (39)$$

from the graphs of Fig. 1d. Note that $\delta m_q^{(\bar{q})} = 0$ because there is no corresponding Feynman graph.

The on-shell renormalization of the squark mixing angle $\theta_{\tilde{q}}$ and $m_q A_q$ is, however, not straightforward. Here we adopt the following procedure: We start from squark pole masses $m_{\tilde{q}_i}$ and the on-shell mixing angle $\theta_{\tilde{q}}$ which will be defined later. The other on-shell parameters $(M_{\tilde{Q},\tilde{U},\tilde{D}}, A_q)$ for squarks are then defined in terms of the above $m_{\tilde{q}_i}$ and $\theta_{\tilde{q}}$ by the tree-level relations eqs. (1)–(5). In this scheme $\delta(m_q A_q)$ takes the form

$$\delta(m_q A_q) = \frac{1}{2}(\delta m_{\tilde{q}_1}^2 - \delta m_{\tilde{q}_2}^2) \sin 2\theta_{\tilde{q}} + (m_{\tilde{q}_1}^2 - m_{\tilde{q}_2}^2) \cos 2\theta_{\tilde{q}} \delta\theta_{\tilde{q}} + \delta m_q \mu \{\cot \beta, \tan \beta\}, \quad (40)$$

where $\cot \beta$ ($\tan \beta$) is for $\tilde{q} = \tilde{t}$ ($\tilde{b}$). Here one has $\delta m_{\tilde{q}_i}^2 = \text{Re}[\Pi_{ii}^{\tilde{q}(g)}(m_{\tilde{q}_i}^2) + \Pi_{ii}^{\tilde{q}(\tilde{g})}(m_{\tilde{q}_i}^2) + \Pi_{ii}^{\tilde{q}(\bar{q})}(m_{\tilde{q}_i}^2)]$ from eqs. (25)–(27).

Next we have to define the on-shell renormalized squark mixing angle $\theta_{\tilde{q}}$. We treated this problem in [11] in the case of $e^+e^- \rightarrow \tilde{q}_i \bar{\tilde{q}}_j$. We fixed the counterterm of the mixing angle $\delta\theta_{\tilde{q}}$ such that it cancels the off-diagonal part of the squark wave-function corrections to $e^+e^- \rightarrow \tilde{q}_1 \bar{\tilde{q}}_2$. Here we use the same scheme and take $\delta\theta_{\tilde{q}} = \delta\theta_{\tilde{q}}^{(\tilde{g})} + \delta\theta_{\tilde{q}}^{(\bar{q})}$ from [11]:

$$\delta\theta_{\tilde{q}}^{(\tilde{g})} = \frac{\alpha_s C_F}{4\pi} \frac{m_{\tilde{g}} m_q}{I_q^{3L}(m_{\tilde{q}_1}^2 - m_{\tilde{q}_2}^2)} (B_0(m_{\tilde{q}_2}^2, m_{\tilde{g}}^2, m_q^2) a_{11} - B_0(m_{\tilde{q}_1}^2, m_{\tilde{g}}^2, m_q^2) a_{22}), \quad (41)$$

$$\delta\theta_{\tilde{q}}^{(\bar{q})} = \frac{\alpha_s C_F}{8\pi} \frac{\sin 4\theta_{\tilde{q}}}{m_{\tilde{q}_1}^2 - m_{\tilde{q}_2}^2} (A_0(m_{\tilde{q}_2}^2) - A_0(m_{\tilde{q}_1}^2)), \quad (42)$$

with $a_{11} = 4(I_q^{3L} \cos^2 \theta_{\tilde{q}} - s_W^2 e_q)$ and $a_{22} = 4(I_q^{3L} \sin^2 \theta_{\tilde{q}} - s_W^2 e_q)$. The counterterms $\delta G_{ijk}^{\tilde{q}(0)}$ are then completely fixed.

In our calculation we need the on-shell parameters $(M_{\tilde{Q},\tilde{U},\tilde{D}}, A_q)$. In a combined treatment of both the stop and the sbottom sectors in the on-shell scheme we have to pay special attention to the parameter $M_{\tilde{Q}}$. This is necessary for the calculation of the

decay width of $H^+ \rightarrow t\bar{b}$ and of the branching ratios of Higgs decays. At tree-level and in the $\overline{\text{DR}}$ scheme the parameter $M_{\tilde{Q}}$ in the stop and sbottom mass matrices must be equal because of $\text{SU}(2)_L$ symmetry. In the on-shell scheme, however, this is not the case. The shifts from the $\overline{\text{DR}}$ parameters to the on-shell (i. e. physical) ones are different for the stop and sbottom sectors:

$$M_{\tilde{Q}}^2|_{\overline{\text{DR}}} = M_{\tilde{Q}}^2(\tilde{t})|_{\text{os}} + \delta M_{\tilde{Q}}^2(\tilde{t}), \quad M_{\tilde{Q}}^2|_{\overline{\text{DR}}} = M_{\tilde{Q}}^2(\tilde{b})|_{\text{os}} + \delta M_{\tilde{Q}}^2(\tilde{b}), \quad (43)$$

with

$$\delta M_{\tilde{Q}}^2(\tilde{q}) = \delta m_{\tilde{q}_1}^2 \cos^2 \theta_{\tilde{q}} + \delta m_{\tilde{q}_2}^2 \sin^2 \theta_{\tilde{q}} - (m_{\tilde{q}_1}^2 - m_{\tilde{q}_2}^2) \sin 2\theta_{\tilde{q}} \delta \theta_{\tilde{q}} - 2m_q \delta m_q. \quad (44)$$

In this paper we take $M_{\tilde{Q}}^2(\tilde{t})|_{\text{os}}$ as the on-shell input parameter. This then leads to a shift of $M_{\tilde{Q}}^2$ in the sbottom sector:

$$M_{\tilde{Q}}^2(\tilde{b})|_{\text{os}} = M_{\tilde{Q}}^2(\tilde{t})|_{\text{os}} + \delta M_{\tilde{Q}}^2(\tilde{t}) - \delta M_{\tilde{Q}}^2(\tilde{b}). \quad (45)$$

As all physical parameters are finite, the shift $\delta M_{\tilde{Q}}^2(\tilde{t}) - \delta M_{\tilde{Q}}^2(\tilde{b})$ has to be UV convergent. We have checked that this is indeed the case.

The one-loop corrected decay width to $\mathcal{O}(\alpha_s)$ in the on-shell scheme is then given by

$$\Gamma(H^k \rightarrow \tilde{q}_i \tilde{\bar{q}}_j) = \frac{N_C \kappa}{16\pi m_{H^k}^3} [|G_{ijk}^{\tilde{q}}|^2 + 2G_{ijk}^{\tilde{q}} \text{Re}(\delta G_{ijk}^{\tilde{q}(v)} + \delta G_{ijk}^{\tilde{q}(w)} + \delta G_{ijk}^{\tilde{q}(0)})]. \quad (46)$$

We have checked the UV convergence of the amplitudes $G_{ijk}^{\tilde{q}\text{corr}}$ of eq. (13) and hence also of $\Gamma(H^k \rightarrow \tilde{q}_i \tilde{\bar{q}}_j)$. The width of eq. (46) is still infrared divergent.

The infrared divergences in (46) are cancelled by including the $\mathcal{O}(\alpha_s)$ contribution from real gluon emission from $\tilde{q}_i$ and $\tilde{\bar{q}}_j$ (see Fig. 1e). The decay width of $H^k(p) \rightarrow \tilde{q}_i(k_1) + \tilde{\bar{q}}_j(k_2) + g(k_3)$ is given by

$$\Gamma(H^k \rightarrow \tilde{q}_i \tilde{\bar{q}}_j g) = \frac{\alpha_s C_F N_C |G_{ijk}^{\tilde{q}}|^2}{4\pi^2 m_{H^k}} [(m_{H^k}^2 - m_{\tilde{q}_i}^2 - m_{\tilde{q}_j}^2) I_{12} - m_{\tilde{q}_i}^2 I_{11} - m_{\tilde{q}_j}^2 I_{22} - I_1 - I_2]. \quad (47)$$

The functions I_n , and I_{nm} are defined as [18]

$$I_{i_1 \dots i_n} = \frac{1}{\pi^2} \int \frac{d^3 k_1}{2E_1} \frac{d^3 k_2}{2E_2} \frac{d^3 k_3}{2E_3} \delta^4(p - k_1 - k_2 - k_3) \frac{1}{(2k_3 k_{i_1} + \lambda^2) \dots (2k_3 k_{i_n} + \lambda^2)}. \quad (48)$$

The explicit forms of $I_{i_1 \dots i_n}$ are given in [18]. In (47), $I_{11,22,12}$ are infrared divergent. We have checked that the infrared divergences in (47) cancel those in (46). In the numerical analysis we define the corrected decay width as

$$\Gamma^{\text{corr}}(H^k \rightarrow \tilde{q}_i \tilde{\bar{q}}_j) \equiv \Gamma(H^k \rightarrow \tilde{q}_i \tilde{\bar{q}}_j) + \Gamma(H^k \rightarrow \tilde{q}_i \tilde{\bar{q}}_j g). \quad (49)$$

4 Numerical results and conclusions

We choose $\{m_{A^0}, m_{t,b}, M, \mu, \tan\beta, M_{\tilde{Q}}(\tilde{t}), A\}$ (with $M_{\tilde{Q}}(\tilde{t}) \equiv M_{\tilde{Q}}(\tilde{t})|_{\text{os}}$) as the basic input parameters of the MSSM, taking $M = (\alpha_2/\alpha_s(m_{\tilde{g}}))m_{\tilde{g}} = (3/5 \tan^2 \theta_W)M'$, $M_{\tilde{Q}}(\tilde{t}) : M_{\tilde{U}} : M_{\tilde{D}} : M_{\tilde{L}} : M_{\tilde{E}} = 1 : \frac{8}{9} : \frac{10}{9} : 1 : 1$ and $A \equiv A_t = A_b = A_\tau$. Here M (M') is the SU(2) (U(1)) gaugino mass, $\alpha_2 = g^2/4\pi$, and $(M_{\tilde{L},\tilde{E}}, A_\tau)$ are the mass matrix parameters of the slepton sector [6, 8]. We take $m_t = 175$ GeV, $m_b = 5$ GeV, $m_Z = 91.2$ GeV, $m_W = 80$ GeV, $\sin^2 \theta_W = 0.23$, $\alpha_2 = 0.0337$, and $\alpha_s = \alpha_s(m_{H^k})$ for the H^k decay. We use $\alpha_s(Q) = 12\pi/\{(33 - 2n_f) \ln(Q^2/\Lambda_{n_f}^2)\}$, with $\alpha_s(m_Z) = 0.12$, and the number of quark flavors $n_f = 5(6)$ for $m_b < Q \leq m_t$ (for $Q > m_t$).

We define the QCD corrections as the difference between the $\mathcal{O}(\alpha_s)$ corrected width $\Gamma^{\text{corr}}(H^k \rightarrow \tilde{q}_i \tilde{\bar{q}}_j)$ of eq. (49) (i. e. eqs. (46) plus (47)) and the tree-level width $\Gamma^{\text{tree}}(H^k \rightarrow \tilde{q}_i \tilde{\bar{q}}_j)$ of eq. (6) with $M_{\tilde{Q}} = M_{\tilde{Q}}(\tilde{t})$ for both the $\tilde{t}$ and $\tilde{b}$ mass matrices. Note that $m_{\tilde{t}_i} = m_{\tilde{t}_i}^{\text{tree}}$ but $m_{\tilde{b}_i} \neq m_{\tilde{b}_i}^{\text{tree}}$ due to the shift in $M_{\tilde{Q}}(\tilde{b})$ in eq. (45), where $m_{\tilde{q}_i}$ ($m_{\tilde{q}_i}^{\text{tree}}$) is the on-shell $\tilde{q}_i$ -mass at one-loop level (at tree-level). This shift is calculated by taking α_s at the scale $M_{\tilde{Q}}(\tilde{t})$ in eq. (44).

In order not to vary too many parameters, in the following we take the values of M, μ , and $\tan\beta$ such that $m_{\tilde{\chi}_1^0} \simeq 70$ GeV as in [8], where $\tilde{\chi}_1^0$ is the lightest neutralino. In Fig. 2 we show the m_{A^0} dependence of the tree-level and corrected widths $\Gamma_{H^k}^{\text{tree}}(\tilde{q}\tilde{\bar{q}}) \equiv \sum_{i,j=1,2} \Gamma^{\text{tree}}(H^k \rightarrow \tilde{q}_i \tilde{\bar{q}}_j)$ and $\Gamma_{H^k}^{\text{corr}}(\tilde{q}\tilde{\bar{q}}) \equiv \sum_{i,j=1,2} \Gamma^{\text{corr}}(H^k \rightarrow \tilde{q}_i \tilde{\bar{q}}_j)$, and the tree-level branching ratio $B_{H^k}^{\text{tree}}(\tilde{q}\tilde{\bar{q}}) \equiv \sum_{i,j=1,2} B^{\text{tree}}(H^k \rightarrow \tilde{q}_i \tilde{\bar{q}}_j)$ [6, 8] for $(\tan\beta, M \text{ (GeV)}, \mu \text{ (GeV)}, M_{\tilde{Q}}(\tilde{t}) \text{ (GeV)}, A \text{ (GeV)}) = (2, 160, 300, 95, 300)$ (a, b, c), and $(12, 140, -300, 150, -250)$ (d, e, f). In these cases we have (in GeV units): $(m_{\tilde{t}_1}, m_{\tilde{t}_2}, m_{\tilde{b}_1}, m_{\tilde{b}_2}, m_{\tilde{b}_1}^{\text{tree}}, m_{\tilde{b}_2}^{\text{tree}}, m_{\tilde{g}}, m_{\tilde{\chi}_1^+}) = (106, 252, 105, 124, 99, 113, 465, 128)$ (a,b,c) and $(97, 297, 113, 215, 102, 210, 412, 133)$ (d,e,f). Here $\tilde{\chi}_1^+$ is the lighter chargino. We see that in these cases the $\tilde{t}\tilde{b}$ mode (the sum of the $\tilde{t}\tilde{t}$ and $\tilde{b}\tilde{b}$ modes) dominates the H^+ decay (the H^0 and A^0 decays) in a wide m_{A^0} range at the tree-level, and that the QCD corrections to the $\tilde{t}\tilde{b}$ mode (the $\tilde{t}\tilde{t}$ and $\tilde{b}\tilde{b}$ modes) are significant, but that as a whole they do not invalidate the dominance of the $\tilde{t}\tilde{b}$ mode (the sum of the $\tilde{t}\tilde{t}$ and $\tilde{b}\tilde{b}$ modes). Our calculation includes the leading Yukawa corrections to the Higgs sector as in [6, 8]. Note that $m_{H^+} \simeq m_{H^0} \simeq m_{A^0}$ in the m_{A^0} range shown here, and that the A^0 does not couple to $\tilde{t}_i \tilde{\bar{t}}_i$ and $\tilde{b}_i \tilde{\bar{b}}_i$ ($i = 1, 2$). As for h^0 decay, we have found that the decay $h^0 \rightarrow \tilde{t}_1 \tilde{\bar{t}}_1$ is kinematically allowed only in a very limited region of the MSSM parameter space [8].

In Table 1 we show the values of the tree-level branching ratios $B^{\text{tree}}(H^+ \rightarrow \tilde{t}\tilde{b})$ (a, d), $B^{\text{tree}}(H^0 \rightarrow \tilde{t}\tilde{t}, \tilde{b}\tilde{b})$ (b, e), $B^{\text{tree}}(A^0 \rightarrow \tilde{t}\tilde{t}, \tilde{b}\tilde{b})$ (c, f), and the QCD corrections $C(H^+ \rightarrow \tilde{t}\tilde{b})$ (a, d), $C(H^0 \rightarrow \tilde{t}\tilde{t}, \tilde{b}\tilde{b})$ (b, e), $C(A^0 \rightarrow \tilde{t}\tilde{t}, \tilde{b}\tilde{b})$ (c, f) for typical values of

$M_{\tilde{Q}}(\tilde{t})$ and A , for $(m_{A^0} \text{ (GeV)}, \tan \beta, \mu \text{ (GeV)}, M \text{ (GeV)}) = (400, 2, 300, 160)$ (a), $(450, 2, 300, 160)$ (b), $(500, 2, 300, 160)$ (c), $(400, 12, -300, 140)$ (d), $(450, 12, -300, 140)$ (e), and $(500, 12, -300, 140)$ (f). Here $B^{\text{tree}}(H^+ \rightarrow \tilde{t}\tilde{b}) \equiv B_{H^+}^{\text{tree}}(\tilde{t}\tilde{b})$, $B^{\text{tree}}(H^k \rightarrow \tilde{t}\tilde{t}, \tilde{b}\tilde{b}) \equiv \sum_{\tilde{q}=\tilde{t},\tilde{b}} B_{H^k}^{\text{tree}}(\tilde{q}\tilde{q})$ ($k = 2, 3$), $C(H^+ \rightarrow \tilde{t}\tilde{b}) \equiv (\Gamma_{H^+}^{\text{corr}}(\tilde{t}\tilde{b}) - \Gamma_{H^+}^{\text{tree}}(\tilde{t}\tilde{b}))/\Gamma_{H^+}^{\text{tree}}(\tilde{t}\tilde{b})$, and $C(H^k \rightarrow \tilde{t}\tilde{t}, \tilde{b}\tilde{b}) \equiv (\sum_{\tilde{q}=\tilde{t},\tilde{b}} \Gamma_{H^k}^{\text{corr}}(\tilde{q}\tilde{q}) - \sum_{\tilde{q}=\tilde{t},\tilde{b}} \Gamma_{H^k}^{\text{tree}}(\tilde{q}\tilde{q}))/\sum_{\tilde{q}=\tilde{t},\tilde{b}} \Gamma_{H^k}^{\text{tree}}(\tilde{q}\tilde{q})$ ($k = 2, 3$). We see again that the QCD corrections are significant, but that the $\tilde{q}\tilde{q}$ modes dominate the H^+ , H^0 , and A^0 decays in a wide region also when the QCD corrections are included. We have found that our results are rather insensitive to the assumptions on the ratios of $M_{\tilde{U},\tilde{D},\tilde{L},\tilde{E}}/M_{\tilde{Q}}(\tilde{t})$ and $A_{b,\tau}/A_t$.

Here we note that for large $\tan \beta$ (and large $|\mu|$) we often get negative corrected widths for some of the $\tilde{b}$ -involved modes (i. e. the $\tilde{b}_i\tilde{b}_j$ and $\tilde{t}_i\tilde{b}_j$ modes) depending on the values of the other input parameters. This is mainly due to a large value of the third term of $\delta(A_b m_b)$ of eq. (40) for large $\tan \beta$ which leads to large values of the shifts $\delta G_{ijk}^{\tilde{b}(0)}$, ($k = 2, 3$) and $\delta G_{ij4}^{(0)}$ of eqs. (32), (36), and (37), which then can result in negative corrected widths for some of the $\tilde{b}$ -involved modes. Here note that the shifts $\delta G_{ijk}^{\tilde{b}(0)}$ and $\delta G_{ij4}^{(0)}$ are roughly proportional to $\delta(A_b m_b) \tan \beta \sim \delta m_b \mu \tan^2 \beta$.

In conclusion, we have calculated the $\mathcal{O}(\alpha_s)$ QCD corrections to the decay widths of $H^+ \rightarrow \tilde{t}\tilde{b}$ and $H^0, A^0 \rightarrow \tilde{t}\tilde{t}, \tilde{b}\tilde{b}$ in the on-shell scheme, including all quark mass terms and $\tilde{q}_L$ - $\tilde{q}_R$ mixing. We find that the QCD corrections are significant, but that they do not invalidate our previous conclusions at tree-level about the dominance of the $\tilde{t}\tilde{b}$, and $\tilde{t}\tilde{t}, \tilde{b}\tilde{b}$ modes in a wide MSSM parameter region.

Acknowledgements

The work of A. B., H. E., and W. M. was supported by the ‘‘Fonds zur F6rderung der wissenschaftlichen Forschung’’ of Austria, project no. P10843-PHY.

References

- [1] See e. g., H. E. Haber and G. L. Kane, Phys. Rep. 117 (1985) 75.
- [2] J. F. Gunion, H. E. Haber, G. L. Kane, and S. Dawson, The Higgs Hunter's Guide, Addison-Wesley (1990).
- [3] J. F. Gunion and H. E. Haber, Nucl. Phys. B272 (1986) 1; B402 (1993) 567 (Erratum).
- [4] Z. Kunszt and F. Zwirner, Nucl. Phys. B385 (1992) 3.
- [5] J. F. Gunion and H. E. Haber, Nucl. Phys. B307 (1988) 445;
A. Djouadi, P. Janot, J. Kalinowski, and P. M. Zerwas, Phys. Lett. B376 (1996) 220.
- [6] A. Bartl, K. Hidaka, Y. Kizukuri, T. Kon, and W. Majerotto, Phys. Lett. B315 (1993) 360.
- [7] A. Djouadi, J. Kalinowski, P. Ohmann, and P. M. Zerwas, DESY-95-213, hep-ph/9605339.
- [8] A. Bartl, H. Eberl, K. Hidaka, T. Kon, W. Majerotto, and Y. Yamada, Phys. Lett. B389 (1996) 538.
- [9] R. A. Jiménez and J. Solà, Phys. Lett. B389 (1996) 53;
J. A. Coarasa, R. A. Jiménez, and J. Solà, Phys. Lett. B389 (1996) 312;
A. Bartl, H. Eberl, K. Hidaka, T. Kon, W. Majerotto, and Y. Yamada, Phys. Lett. B378 (1996) 167.
- [10] A. Bartl, H. Eberl, K. Hidaka, T. Kon, W. Majerotto, and Y. Yamada, Phys. Lett. B373 (1996) 117.
- [11] H. Eberl, A. Bartl, and W. Majerotto, Nucl. Phys. B472 (1996) 481.
- [12] S. Kraml, H. Eberl, A. Bartl, W. Majerotto, and W. Porod, Phys. Lett. B386 (1996) 175.
- [13] A. Djouadi, W. Hollik, and C. Jünger, Phys. Rev. D54 (1996) 5629;
A. Djouadi, W. Hollik, and C. Jünger, Karlsruhe preprint, hep-ph/9609419.
- [14] C. S. Li, R. J. Oakes, and J. M. Yang, Phys. Rev. D54 (1996) 6883.
- [15] W. Beenakker, R. Höpker, T. Plehn, and P. M. Zerwas, DESY 96-178, hep-ph/9610313.

- [16] W. Siegel, Phys. Lett. 84B (1979) 193;
D. M. Capper, D. R. T. Jones and P. van Nieuwenhuizen, Nucl. Phys. B167
(1980) 479;
I. Antoniadis, C. Kounnas and K. Tamvakis, Phys. Lett. 119B (1982) 377.
- [17] I. Jack, D. R. T. Jones, S. P. Martin, M. T. Vaughn and Y. Yamada, Phys. Rev. D50 (1994) 5481.
- [18] A. Denner, Fortschr. Phys. 41 (1993) 307.

Figure Captions

Fig. 1 All diagrams relevant for the calculation of the $\mathcal{O}(\alpha_s)$ QCD corrections to the width of $H^k \rightarrow \tilde{q}_i \tilde{q}_j$ in the MSSM.

Fig. 2 The m_{A^0} dependence of $\Gamma_{H^k}^{\text{tree}}(\tilde{q}\tilde{q})$ (dashed line), $\Gamma_{H^k}^{\text{corr}}(\tilde{q}\tilde{q})$ (solid line), and $B_{H^k}^{\text{tree}}(\tilde{q}\tilde{q})$ (short-dashed line) for $(\tan \beta, M \text{ (GeV)}, \mu \text{ (GeV)}, M_{\tilde{Q}}(\tilde{t}) \text{ (GeV)}, A \text{ (GeV)}) = (2, 160, 300, 95, 300)$ (a, b, c), and $(12, 140, -300, 150, -250)$ (d, e, f).

Table Caption

Table 1 B^{tree} and C for typical values of $M_{\tilde{Q}}(\tilde{t})$ and A , for various values of $(m_{A^0}, \tan \beta, \mu, M)$. See the text for details. The requirement $m_{\tilde{t}_1, \tilde{b}_1, \tilde{\tau}^-} > m_{\tilde{\chi}_1^0} (\simeq 70 \text{ GeV})$ is satisfied for these parameter values.




Fig. 1a




Fig. 1b




Fig. 1c




Fig. 1d




Fig. 1e




Fig.2

Table 1

	$M_{\tilde{Q}}(t)(\text{GeV})$	A(GeV)	B^{tree}	C		$M_{\tilde{Q}}(t)(\text{GeV})$	A(GeV)	B^{tree}	C
(a)	80	0	0.704	-0.002	(d)	140	-250	0.781	0.158
	80	250	0.803	0.125		140	0	0.735	0.121
	120	150	0.732	0.148		140	200	0.756	0.091
	120	350	0.751	-0.052		180	-300	0.680	0.114
	140	0	0.609	0.002		180	0	0.624	0.196
	160	350	0.678	0.185		180	250	0.646	0.159
	180	-150	0.506	-0.103		220	-400	0.669	0.103
	240	550	0.706	0.241		220	350	0.600	0.135
(b)	80	0	0.818	-0.072	(e)	140	-250	0.595	0.201
	140	100	0.752	0.164		180	200	0.560	0.116
	200	-150	0.746	-0.132		200	250	0.564	0.155
	200	0	0.706	0.131		240	-400	0.569	0.018
	200	400	0.644	0.220		260	400	0.536	0.092
	260	-300	0.649	-0.247		300	-600	0.611	0.011
	260	600	0.777	0.153		340	650	0.497	0.075
	360	900	0.865	0.171		400	-950	0.649	0.005
(c)	80	200	0.617	0.135	(f)	140	-250	0.591	0.270
	80	300	0.680	0.118		140	-50	0.503	0.350
	100	300	0.668	0.139		160	0	0.445	0.389
	120	250	0.616	0.185		170	150	0.440	0.318
	140	200	0.550	0.244		180	-200	0.464	0.302
	140	350	0.661	0.186		180	200	0.433	0.345
	180	300	0.505	0.451		200	-350	0.531	0.071
	180	450	0.672	0.221		200	300	0.434	0.402